Journal of Surgical Oncology

Table of Contents

Volume 92, Number 1, October 1, 2005

Guest Editorial: The Global Burden of Cancer
Stephen F. Sener and Nathan Grey 1

Contralateral Uptake and Metastases in Sentinel Lymph Node Mapping for Recurrent Breast Cancer
Anurag Agarwal, Dwight E. Heron, Jules Sumkin, and Jeff Falk 4
Some patients may benefit from sentinel lymph node mapping prior to salvage mastectomy. Identifying uptake in a contralateral sentinel lymph node may change the multi-disciplinary management of recurrent invasive breast cancer to include a contralateral axillary dissection, chemotherapy, and/or RT to the contralateral axilla.

Added Cancer Yield of MRI in Screening the Contralateral Breast of Women Recently Diagnosed With Breast Cancer: Results From the International Breast Magnetic Resonance Consortium (IBMC) Trial
Constance D. Lehman, Jeffrey D. Blume, David Thickman, David A. Bluemke, Etta Pisano, Christiane Kuhl, Thomas B. Julian, Nola Hylton, Paul Weatherall, Michael O'Loughlin, Stuart J. Schnitt, Constantine Gatsonis, and Mitchell D. Schnall 9
Multiple prior studies have reported that MRI can detect disease that is occult on mammography, ultrasound, and physical exam. In women with a recent breast cancer diagnosis, approximately 4% will have an otherwise occult invasive breast cancer detected in the opposite breast by MRI alone.

Commentary
Kirby I. Bland 15

Subjective Morbidity and Quality of Life After Sentinel Node Biopsy and Axillary Lymph Node Dissection for Breast Cancer
Emmanuel Barranger, Gil Dubernard, José Fleurence, Martine Antoine, Emile Darai, and Serge Uzan 17
Sentinel node biopsy is associated with significantly lower mid-term morbidity than axillary dissection.

The Consequences of Long-Time Arm Morbidity in Node-Negative Breast Cancer Patients with Sentinel Node Biopsy or Axillary Clearance
M. Leidenius, M. Leivonen, J. Vironen, and K. von Smitten 23

MRI Detection of Distinct Incidental Cancer in Women With Primary Breast Cancer Studied in IBMC 6883
Mitchell D. Schnall, Jeffery Blume, David A. Bluemke, Gia A. DeAngelis, Nanette DeBruhl, Steven Harms, Sylvia H. Heywang-Köbrunner, Nola Hylton, Christiane K. Kuhl, Etta D. Pisano, Petrina Causer, Stuart J. Schnitt, Stanley F. Smazal, Carol B. Stelling, Constance Lehman, Paul T. Weatherall, and Constantine A. Gatsonis 32

Results of Radiation Therapy Performed After Unplanned Surgery (Without Re-Excision) for Soft Tissue Sarcomas
Lucyna Kepka, Herman D. Suit, Saveli I. Goldberg, Andrew E. Rosenberg, Mark C. Gebhardt, Francis J. Hornicek, and Thomas F. DeLaney 39

Management of Grade III Giant Cell Tumors of Bones
Hsien-Chung Wang, Song-Hsiung Chien, and Gau-Tyan Lin 46

Blood Loss and Transfusion Associated With Musculoskeletal Tumor Surgery
Akira Kawai, Hiroaki Kadota, Umio Yamaguchi, Yuki Morimoto, Toshifumi Ozaki, and Yasuo Beppu 52
We examined the extent of blood loss and the transfusion requirement associated with various kinds of musculoskeletal tumor surgeries in a large, well-defined sample. The median blood loss was 10 ml in biopsies, 100 ml in curettages, 190 ml in amputations or disarticulations, 30 ml in marginal resections (marginal resection for bone tumors, 50 ml; marginal resection for soft tissue tumors, 25 ml), and 380 ml in wide resections (wide resection for bone tumors, 2,155 ml; wide resection for soft tissue tumors, 180 ml).

Tumor Budding and Evidence-Based Treatment of T2 Rectal Carcinomas
Tadahiko Masaki, Hiroyoshi Matsuoka, Masanori Sugiyama, Nobutsugu Abe, Atsuhiko Sakamoto, Toshiaki Watanabe, Hirokazu Nagawa, and Yutaka Atomi 59

Determinants of Unresectability and Outcome of Patients With Occult Colorectal Hepatic Metastases
Joseph J. Bennett, Dincai Cao, and Mitchell C. Posner 64

Clinical Outcome in Patients With Complete Pathologic Response (pT0) to Preoperative Irradiation/Chemo-Irradiation Operated for Locally Advanced or Locally Recurrent Rectal Cancer
Johan N. Wiig, Stein G. Larsen, Svein Dueland, and Karl-Erik Giercksky 70

Preservation of Bowel and Urinary Continence in the Management of Locally Recurrent Rectal Cancer
Nir Wasserberg, Andreas M. Kaiser, Joseph W. Nunoo-Mensah, Peter Biernacki, Thomas Kleisli, and Robert W. Beart, Jr. 76

Volume 92, Number 2, November 1, 2005

Clinicopathological Study of Depth of Subserosal Invasion in Patients With pT2 Gallbladder Carcinoma
Ryoko Sasaki, Noriyuki Uesugi, Hidenori Itabashi, Tomohiro Fujita, Yuichiro Takeda, Koichi Hoshikawa, Masahiro Takahashi, Osamu Funato, Hiroyuki Nitta, Tamotsu Sugai, Senji Kanno, and Kazuyoshi Saito 83
We examined whether depth of subserosal cancer invasion predicts lymph node involvement and survival in gallbladder carcinoma (GBC) patients with pathologicial subserosal invasion (pT2), to explore which patients benefit from radical second resection among patients with inapparent pT2 tumor.

Molecular Alterations During Development of Esophageal Adenocarcinoma
Dorine A. Bax, Peter D. Siersema, Arnoud H.M. Van Vliet, Ernst J. Kuipers, and Johannes G. Kusters 89
Novel insights in the molecular mechanisms that underlie the development of esophageal adenocarcinoma are reviewed, focusing on the role of inflammation, angiogenesis, apoptosis inhibition, loss of cell cycle control, and loss of cell-cell adhesion.

Commentary
Alan G. Casson 99

Significance of the Relationship Between Irregular Regeneration and Two Hepatocarcinogenic Pathways: "De Novo" and So-Called "Dysplastic Nodule-Hepatocellular Carcinoma" Sequence
Kenichi Taguchi, Shinichi Aishima, Shuji Matsuura, Takahiro Terashi, Kenichi Nishiyama, Ken Shirabe, Mitsuo Shimada, Yoshihiko Maehara, and Masazumi Tsuneyoshi 100
Early well differentiated hepatocellular carcinomas are divided into two groups morphologically, with or without dysplastic lesion. Irregular regeneration of hepatocytes impacts on dysplasia-related hepatocarcinogenesis.

Characteristics of Gastric Carcinoma Invading the Muscularis Propria
Eigo Otsuji, Yoshiaki Kuriu, Daisuke Ichikawa, Kazuma Okamoto, Akeo Hagiwara, and Hisakazu Yamagishi 104
Characteristics of gastric carcinomas invading the muscularis propria are compared to carcinomas invading only the mucosa or submucosa, and to those invading the subserosa or serosa. Characteristics of gastric carcinoma invading the muscularis propria are intermediate between those of early and advanced carcinoma.

Deficiency of hMLH1 and hMSH2 Expression Is a Poor Prognostic Factor in Esophageal Squamous Cell Carcinoma
Hirofumi Uehara, Masaki Miyamoto, Kentaro Kato, Yasushi Cho, Takanori Kurokawa, Soichi Murakami, Akira Fukunaga, Yuma Ebihara, Hiroyuki Kaneko, Hiroyuki Hashimoto, Yosihiro Murakami, Toshiaki Shichinohe, You Kawarada, Tomoo Itoh, Shunichi Okushiba, Satoshi Kondo, and Hiroyuki Katoh 109
Loss of MLH1 and/or MSH2 gene expression significantly correlated with increase in malignancy, as evidenced by increase in the existence of metastatic lymph nodes, extensive invasion, and poor differentiation. Patients both MLH1- and MSH2-negative exhibited the poorest clinical outcome compared to other patients.

Reduced p120^{ctn} Expression Correlates With Poor Survival in Patients With Adenocarcinoma of the Gastroesophageal Junction
Bas P.L. Wijnhoven, Massimo Pignatelli, Winand N.M. Dinjens, and Hugo W. Tilanus 116
Expression of p120-catenin, a member of the E-cadherin–catenin cell–cell adhesion complex, was assessed by immunohistochemistry in adenocarcinomas of the gastroesophageal junction. Reduced expression was found in two-third of the tumors and this correlated with a poor 5-year survival.

Surgery Promotes Implantation of Disseminated Tumor Cells, but Does Not Increase Growth of Tumor Cell Clusters
S. Ten Raa, S.J. Oosterling, N.P. van der Kaaij, M.P. van den Tol, R.H.J. Beelen, S. Meijer, C.H.J. van Eijck, J.R.M. van der Sijp, M. van Egmond, and J. Jeekel 124
Effects of local and remote surgery on intra-abdominal tumor development are evaluated.

Pediatric Thyroid Cancer
Devendra A. Chaukar, Venkatesh Rangarajan, Narendra Nair, Anil K. Dcruz, Mandar S. Nadkarni, Prathmesh S. Pai, and Rajesh C. Mistry 130
Thyroid cancer in the pediatric age group presents in an advanced stage with a high incidence of lymph node and distant metastasis. An aggressive approach at the initial surgery is advisable to facilitate postoperative radioiodine scan and is associated with a good long-term prognosis.

Case Report: Forequarter Amputation for Recurrent Breast Cancer: A Case Report and Review of the Literature
Michael D. Goodman, Benjamin McIntyre, Elizabeth A. Shaughnessy, Andrew M. Lowy, and Syed A. Ahmad 134

How I Do It: An Instrument to Provide Containment of Intraoperative Intraperitoneal Chemotherapy With Optimized Distribution
Paul H. Sugarbaker 142

How I Do It: Use of Gold Seeds as Tumor Bed Markers
Subir Nag, J.C. Abdou, and Granger Scruggs 147

Volume 92, Number 3, December 1, 2005

Seminars in Surgical Oncology: Recent Advances in the Management of Esophageal Adenocarcinoma
Alan G. Casson and J. Jan B. van Lanschot, Guest Editors

Preface: Recent Advances in the Management of Esophageal Adenocarcinoma
Alan G. Casson and J. Jan B. van Lanschot 149

Epidemiology of Esophageal Adenocarcinoma
Manuel Pera, Carlos Manterola, Oscar Vidal, and Luis Grande 151

Differences and Similarities of Adenocarcinomas of the Esophagus and Esophagogastric Junction
W.A. Marsman, G.N.J. Tytgat, F.J.W. ten Kate, and J.J.B. van Lanschot 160

The Molecular Biology of Esophageal Adenocarcinoma
Linetta B. Koppert, Bas P.L. Wijnhoven, Herman van Dekken, Hugo W. Tilanus, and Winand N.M. Dinjens 169

Diagnostic Radiology and Nuclear Medicine
Ernesto Castillo and Leo P. Lawler 191

Diagnostic and Therapeutic Endoscopy
E.J. Kuipers and J. Haringsma 203

Surgery for Early Stage Esophageal Adenocarcinoma
H.J. Stein, B.H.A. von Rahden, and M. Feith 210

Surgical Techniques
T. Lerut, W. Coosemans, G. Decker, P. De Leyn, J. Moons, P. Nafteux, and D. Van Raemdonck 218

Neoadjuvant and Adjuvant Chemotherapy for Esophageal Adenocarcinoma
M.M. Mooney 230

Radiation and Chemoradiation Therapy for Esophageal Adenocarcinoma
Jean-François Bosset, F. Lorchel, G. Mantion, J. Buffet, G. Créhange, M. Bosset, L. Chaigneau, and S. Servagi 239

Palliative Therapy
Marjolein Y.V. Homs, Ernst J. Kuipers, and Peter D. Siersema 246

Novel Molecular Targeted Therapy for Esophageal Cancer
David S. Schrump and Dao M. Nguyen 257

Improving Outcomes After Esophagectomy: The Impact of Operative Volume
Alan G. Casson and J. Jan B. van Lanschot 262

Volume 92, Number 4, December 15, 2005

Guest Editorial: cDNA Microarray-Based Translational Research in Soft Tissue Sarcoma
Joanna M. Lubieniecka and Torsten O. Nielsen 267

Guest Editorial: Courageous Challenge: Clinical Intestinal Transplantation
Tokihiko Sawada and Keiichi Kubota 272

Guest Editorial: Predictive Markers for Postsurgical Recurrence of Hepatocellular Carcinoma
Zu-Hua Gao 274

Guest Editorial: Molecular Mechanisms of Hepatic Metastasis in Colorectal Cancer
Heather Rossi 276

Radiochemotherapy With Vinblastine, Methotrexate, and Bleomycin in the Treatment of Verrucous Carcinoma of the Head and Neck
Primož Strojan, Erika Soba, Marjan Budihna, and Marija Auersperg 278
Twelve patients with verrucous carcinoma of the head and neck were treated with radiotherapy and chemotherapy with Vinblastine, Methotrexate, and Bleomycin. Combined radiochemotherapy proved to be highly effective in the treatment of this rare disease with the advantage of organ preservation.

Intestinal Transplantation for the Treatment of Neoplastic Disease
Jang I. Moon, Gennaro Selvaggi, Seigo Nishida, David M. Levi, Tomoaki Kato, Philip Ruiz, Pablo Bejarano, Juan R. Madariaga, and Andreas G. Tzakis 284
Single center experience for the intestinal transplantation to treat intra-abdominal neoplastic disease is reviewed.

Non-Axillary Breast Cancer Recurrences After Sentinel Node Biopsy
Maartje C. van Rijk, Omgo E. Nieweg, Renato A. Valdés Olmos, Emiel J.Th. Rutgers, Cornelis A. Hoefnagel, and Bin B.R. Kroon 292

Detection of Infraclavicular Lymph Node Metastases Using Ultrasonography in Breast Cancer
Huseyln Altinyollar, Gurbuz Dingil, and Ugur Berberoglu 299

Loss of E-cadherin mRNA and Gain of Osteopontin mRNA Are Useful Markers for Detecting Early Recurrence of HCV-Related Hepatocellular Carcinoma
Yukihiro Iso, Tokihiko Sawada, Toshie Okada, and Keiichi Kubota 304

Primary Angiosarcoma of the Spleen
Jun-Te Hsu, Han-Ming Chen, Chin-Yew Lin, Chun-Nan Yeh, Tsann-Long Hwang, Yi-Yin Jan, and Miin-Fu Chen 312

Expression of Angiogenic VEGF-A (Soluble Isoforms 121, 165) and Lymphangiogenic VEGF-C in Colorectal Cancers With Micro-Satellite Instability
Simon J. Hollingsworth, Elizabeth R. Drye, Samson I.H. Tou, and Paul B. Boulos 317
Analysis of VEGF-A and VEGF-C gene activity (transcription and protein expression) suggests that colorectal cancers with high-level micro-satellite instability (MSI-H) have reduced angiogenic/lymphangiogenic potential, helping to explain in part, their better prognosis compared to micro-satellite stable tumours. Transcription of VEGF-121 may be important in the early growth and spread of CRC, and elevated VEGF-121 and VEGF-C transcription with singular di-nucleotide mutations may aid in the identification of distinct MSI-L cancers.

Overexpression of PIM-1 Is a Potential Biomarker in Prostate Carcinoma
Yong Xu, Tong Zhang, Hua Tang, Shumin Zhang, Min Liu, Dalin Ren, and Yuanjie Niu 326

Growth Index, Assessed with Ki-67 and ssDNA Labeling; a Significant Prognosticator for Patients Undergoing Curative Resection for Hepatocellular Carcinoma
Soichiro Morinaga, Naoki Ishiwa, Yoshikazu Noguchi, Yuji Yamamoto, Yasushi Rino, Toshio Imada, Yoshinori Takanashi, Makoto Akaike, Yukio Sugimasa, and Shoji Takemiya 331

Expression of Pro-Apoptotic (p53, p21, bax, bak and fas) and Anti-Apoptotic (bcl-2 and bcl-x) Proteins in Serous Versus Mucinous Borderline Ovarian Tumours
Raffaèle Fauvet, Charlotte Dufournet, Christophe Poncelet, Catherine Uzan, Danièle Hugol, and Emile Daraï 337
p53 overexpression is a common feature of mucinous and serous borderline tumours whereas p21 and bcl-2 overexpression appear specific to serous tumours.

Case Report: Morpheaform Extra-Ocular Sebaceous Carcinoma
D. Innocenzi, A. Balzani, F. Lupi, C. Panetta, N. Skroza, F. Cantoresi, R.A. Schwartz, and S. Calvieri 344

Molecular Mechanisms of Hepatic Metastasis in Colorectal Cancer
L.R. Rudmik and A.M. Magliocco 347
The molecular factors involved in proteolysis, adhesion, angiogenesis, and tumor cell survival in colorectal cancer hepatic metastasis are reviewed. Identification of the molecular mechanisms underlying this complex process could lead to the development of novel screening tools and anti-metastasis therapeutics.

Author Index to Volume 92 361

Subject Index to Volume 92 363